

On the existence of a proper minimal surface in $\mathbb{R}^3$ with the conformal type of a disk.*

Santiago Morales

November 5, 2002

1 Introduction

Proper minimal surfaces in $\mathbb{R}^3$ have peculiar properties that are not shared by general minimal surfaces; especially in the embedded case.

It has been proved that, under additional conditions, this family of surfaces has strong restrictions on their conformal structures. For instance, Huber and Osserman proved that if M is a complete minimal surface with finite total curvature, then M has the conformal type of a compact Riemann surface minus a finite number of points. In particular it is parabolic, that is, M is not compact and M does not carry a negative non-constant subharmonic function.

In the same context, Collin, Kusner, Meeks, and Rosenberg [1] have proved that if M is a properly immersed minimal surface in $\mathbb{R}^3$, then $M(+)=\{(x_1, x_2, x_3) \in M : x_3 \geq 0\}$ is parabolic.

These results have motivated the following conjecture:

Conjecture (Meeks, Sullivan, [3]) *If $f : M \rightarrow \mathbb{R}^3$ is a complete proper minimal immersion where M is a Riemannian surface without boundary and with finite genus, then M is parabolic.*

The main goal of this paper is to show a counterexample to the conjecture.

Theorem *There exists $\chi : \mathbb{D} \rightarrow \mathbb{R}^3$, a conformal proper minimal immersion defined on the unit disk.*

*Research partially supported by DGICYT grant number BFM2001-3489.

2000 Mathematics Subject Classification: Primary 53A10.

Key words and phrases: Minimal Surfaces.

The immersion χ is obtained as the limit of a sequence of minimal immersions with boundary. These boundaries go uniformly to infinity in such a way that we get properness at the limit.

The aforementioned sequence is constructed in a recursive way. In this process, the lemma in page 3 is crucial. In this lemma, we modify a given minimal surface X near its boundary to obtain a new minimal surface Y such that the norm of Y along the boundary increases with respect to the norm of X and, at same time, the norm of Y is controlled by a large enough lower bound in a small neighborhood of the boundary.

The tools we have utilized in the proof of this lemma are those that Nadirashvili [4] used to obtain a complete bounded minimal surface in $\mathbb{R}^3$: Runge's theorem and the López-Ros transformation.

It is important to remark that the geometry of the surface described in the theorem is very complicated; the convex hull of the image under χ of any closed set of $\mathbb{D}$ containing an open arc of $\partial\mathbb{D}$ is $\mathbb{R}^3$.

I would like to point out that the same technique of this paper can be used to construct a complete minimal surface properly immersed in a ball of $\mathbb{R}^3$, [2].

Acknowledgments. I would like to thank Francisco Martín for suggesting to me this line of work and for several informative conversations. I would also like to thank Francisco J. López for helpful criticisms of the paper.

2 Background and Notation

The purpose of this section is to fix the notation used in the paper and to summarize some results about minimal surfaces.

We set $D(z_0, r) = \{z \in \mathbb{C} : |z - z_0| < r\}$. By a *polygon* P we mean a closed simple curve in $\mathbb{R}^2$ formed by a finite number of straight segments verifying $0 \in \text{Int } P$, where $\text{Int } P$ denotes the interior domain bounded by the curve P .

Let $X : D \rightarrow \mathbb{R}^3$ be a conformal minimal immersion defined on a simply connected domain D , and let $S = \{e_1, e_2, e_3\}$ be a set of orthogonal coordinates in $\mathbb{R}^3$. We label $(X(z))_{j,S} = \langle X(z), e_j \rangle$, $j = 1, 2, 3$. We write $(X(z))_j$ instead $(X(z))_{j,S}$ when it is clear which orthogonal frame we are using.

We define the Weierstrass representation of the minimal immersion X in S , $\phi_{(X,S)} = \{\phi_{1,S}, \phi_{2,S}, \phi_{3,S}\}$, as

$$\phi_{j,S}(z) = \frac{\partial(X(z))_{j,S}}{\partial u} - i \frac{\partial(X(z))_{j,S}}{\partial v}, \quad j = 1, 2, 3, \quad z = u + iv,$$

The functions $\phi_{1,S}, \phi_{2,S}, \phi_{3,S}$ are holomorphic on D , verifying $\sum_{j=1}^3 (\phi_{j,S})^2 \equiv 0$ and $\sum_{j=1}^3 |\phi_{j,S}|^2 \not\equiv 0$. As usual, we define:

$$f_{(X,S)} = \phi_{1,S} - i\phi_{2,S} \quad \text{and} \quad g_{(X,S)} = \frac{\phi_{3,S}}{\phi_{1,S} - i\phi_{2,S}}.$$

For ϕ the Weierstrass representation of X , we also denote $f_{(\phi,S)} = f_{(X,S)}$ and $g_{(\phi,S)} = g_{(X,S)}$.

Conversely, if we consider $S = \{e_1, e_2, e_3\}$ a set of Cartesian orthogonal coordinates in $\mathbb{R}^3$, and f and g are (respectively) a holomorphic and meromorphic functions on D such that

$$\phi_{1,S} = \frac{1}{2}f(1 - g^2), \quad \phi_{2,S} = \frac{i}{2}f(1 + g^2), \quad \phi_{3,S} = fg$$

are holomorphic functions on D , then

$$X : D \rightarrow \mathbb{R}^3, \\ X(z) = \sum_{j=1}^3 \left(\operatorname{Re} \int_{z_0}^z \phi_{j,S}(w) dw \right) e_j + c, \quad z_0 \in D, \quad c \in \mathbb{R}^3,$$

is a conformal minimal immersion.

We can write the conformal metric associated to the immersion X , $\lambda_X^2(z) \langle \cdot, \cdot \rangle$, in terms of the Weierstrass representation as follows:

$$\lambda_X(z) = \frac{1}{2} |f_{(X,S)}(z)| (1 + |g_{(X,S)}(z)|^2) = \frac{1}{\sqrt{2}} \|\phi_{(X,S)}(z)\|.$$

Observe that the above formula does not depend on the orthogonal frame S .

3 Proof of the Theorem

In order to prove the theorem, we need the following lemma.

Lemma *Let $X : O \rightarrow \mathbb{R}^3$ be a conformal minimal immersion defined on a simply connected domain O with $X(0) = 0$. Consider $r > 0$, $0 < s < r/100$ and a polygon P with $P \subset O$, satisfying:*

$$r < \|X(z)\| < r + s/2, \quad \forall z \in O \setminus \operatorname{Int} P. \quad (1)$$

Then, for any $b_1, b_2 > 0$, there exist a polygon Q and a conformal minimal immersion $Y : U \rightarrow \mathbb{R}^3$ defined on an open neighborhood of $\overline{\operatorname{Int} Q}$ with $Y(0) = 0$, such that:

- (a) $P \subset \operatorname{Int} Q \subset \overline{\operatorname{Int} Q} \subset U \subset O$;
- (b) $\|Y(z) - X(z)\| < b_1, \forall z \in \overline{\operatorname{Int} P}$;
- (c) $|\|Y(z)\| - (r + s)| < b_2, \forall z \in Q$;
- (d) $\|Y(z)\| > r - 3\sqrt{sr} > r/2, \forall z \in \operatorname{Int} Q \setminus \operatorname{Int} P$.

Roughly speaking, in this lemma, we modify a minimal surface X near the boundary, (property (b)), to obtain a new minimal surface Y such that the norm of Y along a polygon increases with respect to the norm of X , (property (c)), and, at same time, the norm of Y is controlled by a lower bound, (property (d)). Properties (c) and (d) are crucial to obtain properness.

This lemma will be proved in Section 4.

We use the lemma to construct two sequences $\{X_n\}_n$ and $\{P_n\}_n$, where P_n is a polygon, and X_n is a conformal minimal immersion defined on a neighborhood of $\overline{\text{Int } P_n}$ with $X_n(0) = 0$, satisfying the following properties for all $n \in \mathbb{N}$:

$$(\text{T1})_n \quad P_{n-1} \subset \text{Int } P_n \text{ and } P_n \subset D(0, 3);$$

$$(\text{T2})_n \quad |||X_n(z)|| - r_n| < \frac{1}{2(n+1)^2}, \quad \forall z \in P_n;$$

$$(\text{T3})_n \quad ||X_n(z)|| \geq \frac{r_{n-1}}{2} - \frac{1}{2n}, \quad \forall z \in \text{Int } P_n \setminus \text{Int } P_{n-1};$$

$$(\text{T4})_n \quad ||X_n(z) - X_{n-1}(z)|| < \frac{1}{n^2}, \quad \forall z \in \overline{\text{Int } P_{n-1}};$$

$$(\text{T5})_n \quad \lambda_{X_n}(z) \geq \alpha_n \lambda_{X_{n-1}}(z), \quad \forall z \in \overline{\text{Int } P_{n-1}} \text{ where } \{\alpha_k\}_k \text{ is a sequence such that } 0 < \alpha_k < 1 \text{ and } \{\prod_{k=1}^j \alpha_k\}_j \text{ converges to } 1/2.$$

where $r_k = r_{k-1} + 2/k$ for $k > 1$ and $r_1 > 301$.

Sequences $\{X_n\}$ and $\{P_n\}$ are constructed in a recursive way. We can take $X_1(u+iv) = r_1(u, v, 0)$ and $P_1 \subset D(0, 3)$ a suitable polygon satisfying $(\text{T2})_1$. Suppose that we have got $X_1, \dots, X_{n-1}$ and $P_1, \dots, P_{n-1}$.

Now we construct the n^{th} term. We choose $\{\hat{\epsilon}_k\} \searrow 0$, with $\hat{\epsilon}_k < 1/n^2$ for all k . For each k we consider $Y_k : U_k \longrightarrow \mathbb{R}^3$ and Q_k , given by the lemma, for the following data:

$$X = X_{n-1}, \quad P = P_{n-1}, \quad r = r_{n-1} - \frac{1}{n}, \quad s = \frac{3}{n}, \quad b_1 = \hat{\epsilon}_k, \quad b_2 = \frac{1}{2(n+1)^2},$$

and O a simply connected domain with $\overline{\text{Int } P_{n-1}} \subset O \subset D(0, 3)$ and verifying (1). From (b) in the lemma, we deduce that the sequence $\{Y_k\}$ uniformly converges to X_{n-1} on $\overline{\text{Int } P_{n-1}}$. This implies that $\{\lambda_{Y_k}\}$ uniformly converges to $\lambda_{X_{n-1}}$ on $\overline{\text{Int } P_{n-1}}$, and hence there is a $k_0 \in \mathbb{N}$ such that:

$$\lambda_{Y_{k_0}}(z) \geq \alpha_n \lambda_{X_{n-1}}(z), \quad \forall z \in \overline{\text{Int } P_{n-1}}.$$

We define $X_n = Y_{k_0}$, $P_n = Q_{k_0}$. It is easy to check (using the lemma) that X_n and P_n verify $(\text{T1})_n, \dots, (\text{T5})_n$. This concludes the construction of the sequences $\{X_n\}$ and $\{P_n\}$.

Now, we define $\Delta = \bigcup_{n \in \mathbb{N}} \text{Int } P_n$. Δ is a proper simply connected domain of $\mathbb{C}$, (see $(\text{T1})_n$); thus Δ is biholomorphic to a disc.

From $(\text{T4})_n$, we have $\{X_n\}$ is a Cauchy sequence on compact sets in Δ . Then there exists $\chi : \Delta \longrightarrow \mathbb{R}^3$ a harmonic map such that $\{X_n\} \rightarrow \chi$, uniformly on compact sets in Δ . χ has the following properties:

- χ is minimal and conformal.
- χ is an immersion. Indeed, for any $z \in \Delta$ there exists $n_0 \in \mathbb{N}$ such that $z \in \text{Int } P_{n_0}$. From Property (T5)_n one has that,

$$\lambda_{X_k}(z) \geq \alpha_k \lambda_{X_{k-1}}(z) \geq \dots \geq \alpha_k \dots \alpha_{n_0+1} \lambda_{X_{n_0}}(z) \geq \alpha_k \dots \alpha_1 \lambda_{X_{n_0}}(z), \quad \forall k > n_0.$$

Taking the limit as $k \rightarrow \infty$, we deduce that

$$\lambda_\chi(z) \geq \frac{1}{2} \lambda_{X_{n_0}}(z) > 0,$$

and so χ is an immersion.

- χ is proper. Let B be a compact subset of $\mathbb{R}^3$. Consider n_0 such that for all $k > n_0$, one has

$$\|p\| < \frac{r_{k-1}}{2} - \frac{1}{2k} - 2, \quad \forall p \in B \quad (2)$$

and, (using (T3)_n and (T4)_n), for all $z \in \text{Int } P_k \setminus \text{Int } P_{k-1}$:

$$\|\chi(z)\| \geq \frac{r_{k-1}}{2} - \frac{1}{2k} - \|X_k(z) - \chi(z)\| > \frac{r_{k-1}}{2} - \frac{1}{2k} - 2. \quad (3)$$

From (2) and (3), we have $\chi^{-1}(B) \cap (\text{Int } P_k \setminus \text{Int } P_{k-1}) = \emptyset$, $\forall k > n_0$. Then $\chi^{-1}(B) \subset \text{Int } P_{n_0}$. This implies that $\chi^{-1}(B)$ is compact.

This completes the proof of the theorem.

4 Proof of the lemma

To prove the lemma, we modify the immersion X such that it changes almost anything on $\text{Int } P$. The aim of this modification is increasing the norm of X to get (c) and (d) in the lemma at the same time.

The proof of the lemma consists of two inductive process. In the first one, we obtain a new minimal immersion X_n defined on a simply connected domain Ω containing $\overline{\text{Int } P}$. The new immersion X_n is close to X on $\text{Int } P$ and its norm is greater than $r + s$ at $a_1, \dots, a_n$, a finite collection of points around the boundary of Ω .

In the second process, from X_n we obtain a new minimal immersion Y defined on Ω that proves the lemma. To construct Y , we increase the norm of X_n along n curves of $\partial\Omega$ that join the points a_i . As a consequence, Y verify (c) for Q a polygon near $\partial\Omega$. Also, Y is close to X_n on $\text{Int } P$, and then Y and X are close on $\text{Int } P$.

In both processes, we need to control the immersions in such a way that (d) holds.

First, we prepare the first process. We are going to fix some constants and a finite collection of point $p_1, \dots, p_n$ around the polygon P that we will use along this proof. Our purpose is add poles at the points p_i to the Weierstrass representation ϕ^0 of X , in order to increase the norm of the immersion at the aforementioned points $a_1, \dots, a_n$.

We define the following constants:

- 1.a) $\lambda > 1$ such that $\lambda^3 < 2$ and $\lambda(r+s) - 2\lambda^3\sqrt{\lambda^4(r+s)^2 - r^2} > r - 2\sqrt{(r+s)^2 - r^2}$;
- 1.b) $s' = +\sqrt{\lambda^4(r+s)^2 - r^2}$;
- 1.c) $1 > \epsilon_0 > 0$ chosen small enough so that certain inequalities that we are going to use in this paper, were true. ϵ_0 only depends on r, s, b_1, λ and s' .

Let P' be a new polygon and W a simply connected open set such that $P \subset \text{Int } P' \subset \overline{\text{Int } P'} \subset W \subset \overline{W} \subset O$. For a small enough neighborhood E of P' (with $E \subset W \setminus \text{Int } P$), we can define a continuous map S , where $S(z), \forall z \in E$ is a set of orthogonal coordinates in $\mathbb{R}^3$, $\{e_1(z), e_2(z), e_3(z)\}$ with $e_3(z) = \frac{X(z)}{\|X(z)\|}$.




Figure 1: Distribution of the points p_i .

We choose a finite collection of point $\{p_1, \dots, p_n\} \subset E$, (we label $p_{n+1} = p_1$), such that the segments $\overline{p_1 p_2}, \dots, \overline{p_{n-1} p_n}, \overline{p_n p_{n+1}}$ form a new polygon $\hat{P} \subset E$ and verifying the following properties:

2.a) for all $i = 1, \dots, n$, there exists a disc $B^i \subset E$ with $p_i, p_{i+1} \in B^i$ and small enough so that X is close to a constant map on B^i , i.e.

$$\|X(z) - X(w)\| < \epsilon_0, \quad \forall z, w \in B^i; \quad (4)$$

2.b) for all $i = 1, \dots, n$, the sets of orthogonal coordinates $S(p_i)$ and $S(p_{i+1})$ are close, i.e.

$$\|e_j^i - e_j^{i+1}\| < \epsilon_0, \quad \forall j \in \{1, 2, 3\}; \quad (5)$$

where we label $e_j^i = e_j(p_i)$.

2.c) for all $i = 1, \dots, n$,

$$f_{(X, S(p_i))}(p_i) \neq 0. \quad (6)$$

We can choose $p_1, \dots, p_n$ verifying (6) because $f_{(X, S(p))}(p) = 0$ can not hold on any open subset. If we have $f_{(X, S(p))}(p) = 0$ on an open subset then, $X(p)$ is parallel to the normal vector of X at p , $X(p) = \|X(p)\|N(p)$, but this is impossible on an open set for X minimal immersion.

Note that n diverges as $\epsilon_0 \rightarrow 0$.

We define n complex numbers $\theta_1, \dots, \theta_n$, with $|\theta_i| = 1$ and $\text{Im } \theta_i \neq 0$ such that

$$\left| \frac{\overline{f_{(X, S(p_i))}(p_i)}}{|f_{(X, S(p_i))}(p_i)|} - \frac{\overline{f_{(X, S(p_{i+1}))}(p_{i+1})}}{|f_{(X, S(p_{i+1}))}(p_{i+1})|} \right| < \epsilon_0, \quad \forall i = 1, \dots, n. \quad (7)$$

Remark that 2.a), 2.b) and (7) holds for $i = n$ where $p_{n+1} = p_1$.

For every point p_i , $i = 1, \dots, n$, we consider a disk $D(p_i, \delta)$ where $0 < \delta < \epsilon_0^2$ chosen small enough so that, (see Figure 1):

3.a) $\widehat{\text{Int } P} \setminus \cup_{k=1}^n D(p_k, \delta)$ is a simply connected set;

3.b) $\overline{D(p_i, \delta)} \cup \overline{D(p_{i+1}, \delta)} \subset B^i$, $\forall i = 1, \dots, n$;

3.c) $D(p_i, \delta) \cap D(p_k, \delta) = \emptyset$, $\forall i = 1, \dots, n$, and $k \neq i$;

3.d) $\sqrt{\delta} \max_{\overline{D(p_i, \delta)}} \{|f_{(X, S(p_i))}|\} < 1$, $\forall i = 1, \dots, n$;

3.e) $\sqrt{\delta} \frac{\max_{\overline{D(p_i, \delta)}} \{|f_{(X, S(p_i))}|^2 g_{(X, S(p_i))}^2\}}{|\text{Im } \theta_i|} < 1$, $\forall i = 1, \dots, n$;

3.f) $\sqrt{\delta} \max_{\overline{D(p_i, \delta)}} \{\|\phi^0\|\} < 1$, $\forall i = 1, \dots, n$.

We finish these previous steps defining l as

$$l = \sup_{z \in \widehat{\text{Int } P} \setminus \cup_{k=1}^n D(p_k, \delta)} \inf \{ \text{Euclidean length of } \alpha : \alpha \text{ is a curve in } \overline{\widehat{\text{Int } P} \setminus \cup_{k=1}^n D(p_k, \delta)} \text{ with origin } 0 \text{ and ending at } z \} + 2\pi\delta + \delta + 1. \quad (8)$$

Observe that δ depend on ϵ_0 and $\theta_1, \dots, \theta_n$.

4.1 The first inductive process

In this process, we will modify the Weierstrass data of X, ϕ^0 , to produce a sequence of n new Weierstrass data $\phi^1, \dots, \phi^n$. In the i^{th} step, we modify ϕ^{i-1} on the disk $D(p_i, \delta)$ to obtain a new Weierstrass representation ϕ^i with a pole at p_i that is close to ϕ^{i-1} outside some neighborhoods of $p_1, \dots, p_i$, (see next Property (A6) _{i}). To do this we will use the López-Ros transformation in the orthogonal frame $\overline{S(p_i)}$. At the same time, we have to have a control on ϕ^i along the segment $\overline{p_i(p_i + \delta)}$ (see property (A7) _{i} , (A8) _{i} and (A9) _{i}).

When we have $\phi^1, \dots, \phi^n$, we are going to define a simply connected domain Ω that contains $\overline{\text{Int } P}$ and does not contain any poles of $\phi^1, \dots, \phi^n$. Then $X_i(z) = \int_0^z \phi^i$, $z \in \Omega$ is well defined for all $i = 1, \dots, n$. From (A6) _{i} , we have that X_i and X_{i-1} are close outside $D(p_i, \delta)$ (see property (P1) _{i}). In particular X_n are close to X_1 outside $\Omega \setminus \cup_{k=1}^n D(p_k, \delta)$. As ϕ^i has a pole at p_i , we obtain a point a_i that is close to p_i and verifies (A7) _{i} , that is, $\|X_n(a_i)\|$ is greater than $r + s$, (see property (P4) _{i}). The control that we have on ϕ^i along the segment $\overline{p_i(p_i + \delta)}$ implies that $X_n(a_i)$ is close to $X_n(a_{i+1})$ (see property (P3) _{i}). This fact will be crucial along the second inductive process, Section 4.2.

We are going to construct in a recursive way a sequence $\Psi_i = \{\phi^i, k_i, a_i, C_i, G_i, D_i\}$, $i = 1, \dots, n$, where:

- 4.a) $\phi^i : \overline{W} \rightarrow \mathbb{C}^3$ is a Weierstrass representation with poles at $p_1, \dots, p_i$;
- 4.b) k_i is a suitable positive constant;
- 4.c) a_i is a point lying on the segment $\overline{p_i q_i}$, where $q_i = p_i + \delta$;
- 4.d) C_i is an open arc of a circumference centered at p_i with $a_i \in C_i$;
- 4.e) G_i is the closed annular sector bounded by C_i , a piece of $\partial D(p_i, \delta)$ and two radii as Figure 2 indicates;
- 4.f) D_i is an open simply connected subset of $\mathbb{C}$ verifying $\overline{D_i} \cap G_i = \emptyset$ and

$$\{p_i, w_i = p_i - k_i \theta_i\} \subset D_i \subset \overline{D_i} \subset D(p_i, \delta).$$

Remark 1 *In what follows, we will use the convention that $\Psi_{n+1} \stackrel{\text{def}}{=} \Psi_1$.*

The above sequence is constructed in order to satisfy the following properties:

$$(A1)_i \quad \sqrt{\delta} \max_{\overline{D(p_k, \delta)}} \{|f(\phi^i, S(p_k))|\} < 1, \quad k = i + 1, \dots, n;$$




Figure 2: The points q_i and a_i .

$$(A2)_i \quad \sqrt{\delta} \frac{\max_{D(p_k, \delta)} \{|f_{(\phi^i, S(p_k))} g_{(\phi^i, S(p_k))}^2|\}}{|\operatorname{Im} \theta_k|} < 1, \quad k = i+1, \dots, n;$$

$$(A3)_i \quad 2\lambda^2 s' \frac{|f_{(\phi^i, S(p_k))}(p_k) - f_{(\phi^0, S(p_k))}(p_k)|}{|f_{(\phi^0, S(p_k))}(p_k)|} < \epsilon_0, \quad k = i+1, \dots, n;$$

$$(A4)_i \quad \|\operatorname{Re} \int_{\alpha_z} \phi^i\| < \epsilon_0, \quad \forall z \in C_i \text{ where } \alpha_z \text{ is a piece of } C_i \text{ joining } a_i \text{ and } z;$$

$$(A5)_i \quad \text{In the orthogonal frame } S(p_i), \text{ one has } \phi_3^i = \phi_3^{i-1};$$

$$(A6)_i \quad \|\phi^i(z) - \phi^{i-1}(z)\| < \frac{\epsilon_0}{nl}, \quad \forall z \in \overline{W} \setminus (D(p_i, \delta) \cup (\cup_{k=1}^{i-1} D_k));$$

$$(A7)_i \quad \text{Using the frame } S(p_i), \text{ we have } \|(\operatorname{Re} \int_{q_i a_i} \phi_1^i) e_1^i + (\operatorname{Re} \int_{q_i a_i} \phi_2^i) e_2^i\| > \lambda s';$$

$$(A8)_i \quad \text{In } S(p_i), \text{ one has } \|(\operatorname{Re} \int_{q_i z} \phi_1^i) e_1^i + (\operatorname{Re} \int_{q_i z} \phi_2^i) e_2^i\| < \lambda^3 s', \quad \forall z \in G_i;$$

$$(A9)_i \quad \|\operatorname{Re} \int_{q_i a_i} \phi^i - \operatorname{Re} \int_{q_{i-1} a_{i-1}} \phi^{i-1}\| < \epsilon_0(4\lambda^2 s' + 12), \quad i = 2, \dots, n+1.$$

All the above properties are true for $i = 1, \dots, n$, except for Properties $(A1)_i$, $(A2)_i$ and $(A3)_i$ that hold only for $i = 1, \dots, n-1$. In the same way, Property $(A9)_i$ is valid for $i = 2, \dots, n+1$ (see Remark 1).

As we have mentioned at the beginning of this section, we proceed by recursion.

Let ϕ^0 be the Weierstrass representation of $X = X_0$. Assume we have constructed $\Psi_1, \dots, \Psi_{i-1}$. We are going to construct Ψ_i .

Define the meromorphic data ϕ^i , in the orthogonal frame $S(p_i)$, as follows:

$$f_{(\phi^i, S(p_i))} = f_{(\phi^{i-1}, S(p_i))} h_i, \quad g_{(\phi^i, S(p_i))} = g_{(\phi^{i-1}, S(p_i))} / h_i,$$

where

$$h_i(z) = \frac{k_i \theta_i}{z - p_i} + 1.$$

Furthermore, we assume that $k_i > 0$ is small enough to verify:

$$k_i \max_{w \in \overline{D(p_i, \delta)}} \left\{ \left| \frac{f_{(\phi^{i-1}, S(p_i))}(w) - f_{(\phi^{i-1}, S(p_i))}(p_i)}{w - p_i} \right| \right\} < 1, \quad (9)$$

and $(A1)_i$, $(A2)_i$, $(A3)_i$ and $(A6)_i$. It is possible because $\phi^i \rightarrow \phi^{i-1}$, as $k_i \rightarrow 0$, uniformly on $\overline{W} \setminus (D(p_i, \delta) \cup (\cup_{k=1}^{i-1} D_k))$ and $(A1)_{i-1}$, $(A2)_{i-1}$ and $(A3)_{i-1}$ hold. In the case $i = 1$, notice that we have chosen δ such that $\mathcal{B}.d)$ and $\mathcal{B}.e)$ were true.

Remark 2 *The meromorphic function h_i is close to 1 outside a neighborhood of p_i . The constant θ_i has the effect of a rotation over h_i , when z is close to p_i . Outside a neighborhood of p_i this "rotation effect" almost disappears.*

From the definition of ϕ^i , $(A5)_i$ trivially holds.

We define a_i as the first point in $\overline{q_i p_i}$ when we move along this segment from q_i to p_i , and such that:

$$\frac{1}{2} |f_{(\phi^0, S(p_i))}(p_i)| \int_{\overline{q_i a_i}} \frac{k_i dw}{w - p_i} = \lambda^2 s'. \quad (10)$$

Observe that $f_{(\phi^0, S(p_i))}(p_i) \neq 0$ and $\int_{\overline{q_i z}} \frac{k_i dw}{w - p_i} \in \mathbb{R}^+$, $\forall z \in \overline{q_i p_i}$.

Our next step consist of seeing the following:

$$\left\| \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_1^i(w) dw \right) e_1^i + \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_2^i(w) dw \right) e_2^i - \frac{1}{2} \left(\int_{\overline{q_i z}} \frac{k_i dw}{w - p_i} \right) \left(\operatorname{Re} \overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i} e_1^i + \operatorname{Im} \overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i} e_2^i \right) \right\| < 2\epsilon_0, \quad (11)$$

where $z \in \overline{q_i a_i}$, and ϕ_1^i and ϕ_2^i are expressed in the orthogonal frame $S(p_i)$.

In other words, the first two coordinates in $S(p_i)$ of the curve $z \mapsto \operatorname{Re} \int_{\overline{q_i z}} \phi^i(w) dw$, $z \in \overline{q_i a_i}$, approximates the segment starting at 0 in the direction of $\operatorname{Re} \overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i} e_1^i + \operatorname{Im} \overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i} e_2^i$.

To get (11), for $z \in \overline{q_i a_i}$, we write:

$$\begin{aligned} & \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_1^i(w) dw \right) + i \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_2^i(w) dw \right) = \\ & = \frac{1}{2} \left(\int_{\overline{q_i z}} \overline{f_{(\phi^{i-1}, S(p_i))}(w) h_i(w)} dw - \int_{\overline{q_i z}} f_{(\phi^{i-1}, S(p_i))}(w) g_{(\phi^{i-1}, S(p_i))}^2(w) \frac{dw}{h_i(w)} \right) = \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \int_{\overline{q_i z}} \overline{f_{(\phi^0, S(p_i))}(p_i) \frac{k_i \theta_i}{w - p_i}} dw + \\
&+ \frac{1}{2} \int_{\overline{q_i z}} \overline{(f_{(\phi^{i-1}, S(p_i))}(p_i) - f_{(\phi^0, S(p_i))}(p_i)) \frac{k_i \theta_i}{w - p_i}} dw + \tag{12}
\end{aligned}$$

$$+ \frac{1}{2} \int_{\overline{q_i z}} \overline{(f_{(\phi^{i-1}, S(p_i))}(w) - f_{(\phi^{i-1}, S(p_i))}(p_i)) \frac{k_i \theta_i}{w - p_i}} dw + \tag{13}$$

$$+ \frac{1}{2} \int_{\overline{q_i z}} \overline{f_{(\phi^{i-1}, S(p_i))}(w)} dw - \tag{14}$$

$$- \frac{1}{2} \int_{\overline{q_i z}} f_{(\phi^{i-1}, S(p_i))}(w) g_{(\phi^{i-1}, S(p_i))}^2(w) \frac{dw}{h_i(w)} \tag{15}$$

To obtain an upper bound in (12) we use the definition of a_i and Property (A3) _{$i-1$} in the following sense:

$$\begin{aligned}
\left| \int_{\overline{q_i z}} \overline{(f_{(\phi^{i-1}, S(p_i))}(p_i) - f_{(\phi^0, S(p_i))}(p_i)) \frac{k_i \theta_i}{w - p_i}} dw \right| &\leq \\
&\leq 2\lambda^2 s' \frac{|f_{(\phi^{i-1}, S(p_i))}(p_i) - f_{(\phi^0, S(p_i))}(p_i)|}{|f_{(\phi^0, S(p_i))}(p_i)|} < \epsilon_0. \tag{16}
\end{aligned}$$

Now, we deal with (13). Taking (9) into account one has:

$$\begin{aligned}
\left| \int_{\overline{q_i z}} \overline{(f_{(\phi^{i-1}, S(p_i))}(w) - f_{(\phi^{i-1}, S(p_i))}(p_i)) \frac{k_i \theta_i}{w - p_i}} dw \right| &\leq \\
&\leq \delta k_i \max_{w \in D(p_i, \delta)} \left\{ \left| \frac{f_{(\phi^{i-1}, S(p_i))}(w) - f_{(\phi^{i-1}, S(p_i))}(p_i)}{w - p_i} \right| \right\} < \delta < \epsilon_0. \tag{17}
\end{aligned}$$

To get the same upper bound for (14), we use (A1) _{$i-1$} ,

$$\left| \int_{\overline{q_i z}} \overline{f_{(\phi^{i-1}, S(p_i))}(w)} dw \right| \leq \delta \max_{D(p_i, \delta)} \{|f_{(\phi^{i-1}, S(p_i))}(w)|\} < \sqrt{\delta} < \epsilon_0. \tag{18}$$

In order to bound (15), observe that $|h_i(w)| = \sqrt{(\frac{k_i}{w-p_i})^2 + 2 \operatorname{Re} \theta_i (\frac{k_i}{w-p_i}) + 1} > \sqrt{1 - \operatorname{Re}^2 \theta_i} = |\operatorname{Im} \theta_i|$, $\forall w \in \overline{q_i p_i}$. Therefore, (A2) _{$i-1$} leads us to

$$\begin{aligned}
\left| \int_{\overline{q_i z}} f_{(\phi^{i-1}, S(p_i))}(w) g_{(\phi^{i-1}, S(p_i))}^2(w) \frac{dw}{h_i(w)} \right| &\leq \\
&\leq \delta \frac{\max_{D(p_i, \delta)} \{|f_{(\phi^{i-1}, S(p_i))}(w) g_{(\phi^{i-1}, S(p_i))}^2(w)|\}}{|\operatorname{Im} \theta_i|} < \sqrt{\delta} < \epsilon_0. \tag{19}
\end{aligned}$$

Inequalities (16), (17), (18) and (19) give (11).

From (11) and the definition of a_i we deduce $(A7)_i$, and

$$\left\| \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_1^i \right) e_1^i + \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_2^i \right) e_2^i \right\| < \lambda^2 s' + 2\epsilon_0 < \lambda^3 s', \quad \forall z \in \overline{q_i a_i}. \quad (20)$$

At this point, we are able to define D_i , C_i and G_i .

Let D_i be a simply connected domain with $\{p_i, p_i - k_i \theta_i\} \subset D_i \subset \overline{D_i} \subset D(p_i, \delta)$ and $\overline{D_i} \cap \overline{q_i a_i} = \emptyset$ (this is possible because $\operatorname{Im} \theta_i \neq 0$, thus we have $p_i - k_i \theta_i \notin \overline{p_i q_i}$). Since (20), we can take C_i an open arc of a circumference centered at p_i with $a_i \in C_i$ and small enough to verify $(A4)_i$ and $(A8)_i$ on G_i , where G_i is the closed annular sector bounded by C_i , a piece of $\partial D(p_i, \delta)$ and two radii, as it is indicated in Figure 2.

To check $(A9)_i$, we write:

$$\begin{aligned} & \left\| \operatorname{Re} \int_{\overline{q_i a_i}} \phi^i - \operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi^{i-1} \right\| = \\ & = \left\| \sum_{k=1}^3 \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_{k, S(p_i)}^i \right) e_k^i - \sum_{k=1}^3 \left(\operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{k, S(p_{i-1})}^{i-1} \right) e_k^{i-1} \right\| \leq \\ & \leq \sum_{k=1}^3 \left\| \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_{k, S(p_i)}^i \right) e_k^i - \left(\operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{k, S(p_{i-1})}^{i-1} \right) e_k^{i-1} \right\|. \quad (21) \end{aligned}$$

Next step consists of getting an upper bound for the three addends in (21):

From (11), we have

$$\begin{aligned} & \left\| \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_{1, S(p_i)}^i(w) dw \right) e_1^i - \left(\operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{1, S(p_{i-1})}^{i-1}(w) dw \right) e_1^{i-1} \right\| \leq \\ & \leq \left\| \left(\frac{1}{2} \operatorname{Re} \overline{f_{(\phi^0, S(p_i))}(p_i)} \theta_i \int_{\overline{q_i a_i}} \frac{k_i dw}{w - p_i} \right) e_1^i - \right. \\ & \left. - \left(\frac{1}{2} \operatorname{Re} \overline{f_{(\phi^0, S(p_{i-1}))}(p_{i-1})} \theta_{i-1} \int_{\overline{q_{i-1} a_{i-1}}} \frac{k_{i-1} dw}{w - p_{i-1}} \right) e_1^{i-1} \right\| + 4\epsilon_0 = \end{aligned}$$

using (10),

$$\begin{aligned}
&= \left\| \lambda^2 s' \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_i))}(p_i) \theta_i}}{|f_{(\phi^0, S(p_i))}(p_i)|} e_1^i - \lambda^2 s' \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} e_1^{i-1} \right\| + 4\epsilon_0 \leq \\
&\leq \lambda^2 s' \left\| \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_i))}(p_i) \theta_i}}{|f_{(\phi^0, S(p_i))}(p_i)|} e_1^i - \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} e_1^i \right\| + \\
&+ \lambda^2 s' \left\| \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} e_1^i - \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} e_1^{i-1} \right\| + 4\epsilon_0 \leq \\
&\leq \lambda^2 s' \left| \frac{\overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i}}{|f_{(\phi^0, S(p_i))}(p_i)|} - \frac{\overline{f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} \right| + \\
&+ \lambda^2 s' \|e_1^i - e_1^{i-1}\| \left| \frac{\overline{\operatorname{Re} f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}}}{|f_{(\phi^0, S(p_{i-1}))}(p_{i-1})|} \right| + 4\epsilon_0 \leq
\end{aligned}$$

(7) and (5) apply and conclude:

$$\leq 2\lambda^2 s' \epsilon_0 + 4\epsilon_0 = \epsilon_0(2\lambda^2 s' + 4).$$

In the same way, we obtain a bound for the second addends in (21),

$$\begin{aligned}
&\left\| \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_{2, S(p_i)}^i \right) e_2^i - \left(\operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{2, S(p_{i-1})}^{i-1} \right) e_2^{i-1} \right\| \leq \\
&\leq \left\| \left(\frac{1}{2} \operatorname{Im} \overline{f_{(\phi^0, S(p_i))}(p_i) \theta_i} \int_{\overline{q_i a_i}} \frac{k_i dw}{w - p_i} \right) e_2^i - \right. \\
&\left. - \left(\frac{1}{2} \operatorname{Im} \overline{f_{(\phi^0, S(p_{i-1}))}(p_{i-1}) \theta_{i-1}} \int_{\overline{q_{i-1} a_{i-1}}} \frac{k_{i-1} dw}{w - p_{i-1}} \right) e_2^{i-1} \right\| + 4\epsilon_0 \leq \epsilon_0(2\lambda^2 s' + 4).
\end{aligned}$$

To estimate the third addend in (21), we use the Properties (A6)_k, $k = 1, \dots, i-1$, and 3.f),

$$\begin{aligned}
&\left\| \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_{3, S(p_i)}^i \right) e_3^i - \left(\operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{3, S(p_{i-1})}^{i-1} \right) e_3^{i-1} \right\| \leq \\
&\leq \left| \operatorname{Re} \int_{\overline{q_i a_i}} \phi_{3, S(p_i)}^{i-1} \right| + \left| \operatorname{Re} \int_{\overline{q_{i-1} a_{i-1}}} \phi_{3, S(p_{i-1})}^{i-2} \right| \leq \\
&\leq \delta \left(\max_{D(p_i, \delta)} \{\|\phi^{i-1}\|\} + \max_{D(p_{i-1}, \delta)} \{\|\phi^{i-2}\|\} \right) \leq \\
&\leq \delta \left(\max_{D(p_i, \delta)} \{\|\phi^0\|\} + \max_{D(p_{i-1}, \delta)} \{\|\phi^0\|\} + 2\epsilon_0 \right) \leq 2\sqrt{\delta} + 2\delta\epsilon_0 < 4\epsilon_0.
\end{aligned}$$

The above bounds imply (A9)_i.

Note that case $i = n + 1$:

$$\left\| \left(\operatorname{Re} \int_{\overline{q_1 a_1}} \phi_{1,S(p_1)}^1(w) dw \right) e_1^1 - \left(\operatorname{Re} \int_{\overline{q_n a_n}} \phi_{1,S(p_n)}^n(w) dw \right) e_1^n \right\| \leq \epsilon_0(4\lambda^2 s' + 12),$$

can be proved in the same way.

Afterwards, we have just defined a sequence of Weierstrass representation $\phi^0, \phi^1, \dots, \phi^n$ verifying the above properties (A1)_i, ..., (A9)_i.

Now, we are going to take a domain Ω where immersions $X_i(z) \stackrel{\text{def}}{=} \operatorname{Re} \int_0^z \phi^i(w) dw$, $\forall i = 1, \dots, n$ are well defined and then become minimal immersions. Later, we will describe some properties of these immersion that we will use in the second process.

We define Ω as follows, (see figure 3): Let D^i be a disk centered at p_i containing $D(p_i, \delta)$ for all $i = 1, \dots, n$, and let $\alpha_i \subset D^i \setminus \overline{D(p_i, \delta)}$ be a simple curve connecting $\partial D^i \cap \partial(\operatorname{Int} \hat{P} \setminus \bigcup_{k=1}^n D^k)$ to the point q_i . Let N_i be a small neighborhood of $\alpha_i \cup \overline{q_i a_i}$ in $\overline{G_i} \cup (D^i \setminus D(p_i, \delta))$. We define

$$\Omega = \left(\operatorname{Int} \hat{P} \setminus \bigcup_{k=1}^n D^k \right) \cup \left(\bigcup_{k=1}^n N_k \right).$$




Figure 3: The domain Ω .

For suitable D^i , α_i and N_i , the following properties hold:

5.a) $\overline{\Omega}$ is a simply connected domain;

5.b) $\overline{q_i a_i} \subset \overline{\Omega}$ and $\overline{\text{Int } P} \subset \Omega$;

5.c) $\overline{\Omega}$ does not contain any p_i point and any zero of h_i for all $i = 1, \dots, n$;

5.d) $\sup_{z \in \overline{\Omega}} \inf\{\text{Euclidean length of } \alpha : \alpha \text{ is a curve in } \overline{\Omega} \text{ with origin } 0 \text{ and ending at } z\} < l$ where l has been defined on (8);

5.e) $\overline{\Omega} \cap \overline{D(p_i, \delta)} \subset G_i$.

Taking 5.a) and 5.c) into account, we can define n minimal immersions $X_1, \dots, X_n$,

$$X_i : \Omega' \rightarrow \mathbb{R}^3$$

$$X_i(z) = \text{Re} \int_0^z \phi^i(w) dw, \quad i = 1, \dots, n,$$

where Ω' is a suitable open neighborhood of $\overline{\Omega}$.

Consider an orthogonal frame $T_i = \{w_1^i, w_2^i, w_3^i\}$ for every $i = 1, \dots, n$ in such a way that

$$w_3^i = \frac{X_n(a_i)}{\|X_n(a_i)\|}. \quad (22)$$

These orthogonal frames will use in the second process.

We write Q_i , $i = 1, \dots, n$ as simple curves such that:

6.a) Q_i is connected, $\overline{Q_i} = Q_i$ and $Q_i \cap Q_j = \emptyset$ for $j \neq i$;

6.b) Q_i connects C_i with C_{i+1} ;

6.c) $\partial\Omega$ is formed by the curves Q_i and pieces of C_i ;

6.d) $Q_i \subset B^i$;

6.e) $Q_i \cap \overline{D(p_k, \delta)} = \emptyset$ for $k \notin \{i, i+1\}$;

6.f) We can suppose that $f_{(X_n, T_i)}(z) \neq 0$, $\forall z \in Q_i$, $\forall i = 1, \dots, n$. To obtain this, we only need to make slight local modifications on the choice of Ω . Note that the above modifications can be made in such a way that the properties of Ω and Q_i remain.

Now, we see some properties of the immersions $X_1, \dots, X_n$ for $i = 1, \dots, n$:

(P1) _{i} $\|X_i(z) - X_{i-1}(z)\| < \frac{\epsilon_0}{n}$, $\forall z \in \overline{\Omega} \setminus D(p_i, \delta)$;

(P2) _{i} $(X_i)_3 = (X_{i-1})_3$, in the orthogonal frame $S(p_i)$;

(P3) _{i} $\|X_n(a_i) - X_n(a_{i+1})\| < \epsilon_0(4\lambda^2 s' + 17)$. Observe that $a_{n+1} = a_1$ when $i = n$;

$$(\mathbf{P4})_i \quad \|X_n(a_i)\| > \lambda(r + s);$$

$$(\mathbf{P5})_i \quad \|X_n(q_i) - X_n(z)\| < \lambda^3 s' + 5\epsilon_0, \forall z \in \overline{G_i \cap \Omega}.$$

Property $(\mathbf{P1})_i$ is a easy consequence of $(\mathbf{A6})_i$ and 5.d). On the other hand, it is straightforward to check $(\mathbf{P2})_i$ from $(\mathbf{A5})_i$.

Taking into account $(\mathbf{P1})$, $(\mathbf{A9})_{i+1}$ and (4), we get $(\mathbf{P3})_i$ as follows:

$$\begin{aligned} \|X_n(a_i) - X_n(a_{i+1})\| &\leq \|X_i(a_i) - X_{i+1}(a_{i+1})\| + 2\epsilon_0 \leq \\ &\leq \left\| \operatorname{Re} \int_{\overline{q_i a_i}} \phi^i - \operatorname{Re} \int_{\overline{q_{i+1} a_{i+1}}} \phi^{i+1} \right\| + \|X_i(q_i) - X_{i+1}(q_{i+1})\| + 2\epsilon_0 \leq \\ &\leq \epsilon_0(4\lambda^2 s' + 12) + \|X_0(q_i) - X_0(q_{i+1})\| + 4\epsilon_0 \leq \\ &\leq \epsilon_0(4\lambda^2 s' + 12) + 5\epsilon_0 = \epsilon_0(4\lambda^2 s' + 17). \end{aligned}$$

In order to prove $(\mathbf{P4})_i$, recall that

$$\|X_n(a_i)\| > \|X_i(a_i)\| - \epsilon_0.$$

We give bounds for the coordinates of $X_i(a_i)$ in $S(p_i)$ separately. Firstly, we bound the first and second coordinates.

$$\begin{aligned} &\|(X_i(a_i))_{1,S(p_i)} e_1^i + (X_i(a_i))_{2,S(p_i)} e_2^i\| > \\ &> \left\| \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_1^i \right) e_1^i + \left(\operatorname{Re} \int_{\overline{q_i a_i}} \phi_2^i \right) e_2^i \right\| - \|(X_i(q_i))_1 e_1^i + (X_i(q_i))_2 e_2^i\| > \end{aligned}$$

using $(\mathbf{A7})_i$, (4) and the definition of $S(p_i)$ in page 6, we have

$$\begin{aligned} &> \lambda s' - \|(X_0(q_i))_1 e_1^i + (X_0(q_i))_2 e_2^i\| - \epsilon_0 \geq \\ &\geq \lambda s' - \|(X_0(p_i))_1 e_1^i + (X_0(p_i))_2 e_2^i\| - 2\epsilon_0 = \lambda s' - 2\epsilon_0 > s'. \end{aligned}$$

Now, we get a bound for the third coordinate. We use $(\mathbf{P2})_i$ and Hypothesis (1) in the lemma:

$$\begin{aligned} |(X_i(a_i))_3| &= |(X_{i-1}(a_i))_3| \geq |(X_0(a_i))_3| - \epsilon_0 \geq \\ &\geq |(X_0(p_i))_3| - 2\epsilon_0 = \|X_0(p_i)\| - 2\epsilon_0 > r - 2\epsilon_0. \end{aligned}$$

Then, one has

$$\begin{aligned} \|X_n(a_i)\| &> \sqrt{(s')^2 + (r - 2\epsilon_0)^2} - \epsilon_0 = \\ &= \sqrt{\lambda^4(r + s)^2 - 4\epsilon_0(r - \epsilon_0)} - \epsilon_0 > \lambda(r + s), \end{aligned}$$

that prove $(\mathbf{P4})_i$.

To obtain $(P5)_i$ we consider the orthogonal frame $S(p_i)$. Let $z \in \overline{G_i \cap \Omega}$. Using $(A8)_i$, we have

$$\begin{aligned} \|X_n(q_i) - X_n(z)\| &\leq \|X_i(q_i) - X_i(z)\| + 2\epsilon_0 \leq \\ &\leq \left\| \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_1^i \right) e_1^i + \left(\operatorname{Re} \int_{\overline{q_i z}} \phi_2^i \right) e_2^i \right\| + |(X_i(q_i) - X_i(z))_{3,S(p_i)}| + 2\epsilon_0 < \\ &< \lambda^3 s' + |(X_0(q_i) - X_0(z))_{3,S(p_i)}| + 4\epsilon_0 \leq \lambda^3 s' + 5\epsilon_0. \end{aligned}$$

4.2 The second inductive process

In the first process, we have obtained a immersion X_n ; and we have only guaranteed that its norm is greater than $r + s$ at the points $a_1, \dots, a_n$, (Property (P4)). In this second process, our objective is increasing the norm of X_n around the whole boundary of Ω in order to satisfy its norm is greater than $r + s$. So, we will modify the immersion X_n to produce a sequence of minimal immersions $Y_1, \dots, Y_n$ defined on $\overline{\Omega}$. In each step, we take certain small neighborhoods, Q_i^ξ of the curve Q_i ; and modify the Weierstrass representation of Y_{i-1} , ψ^{i-1} , on Q_i^ξ , to get a new minimal immersion Y_i . Y_i and Y_{i-1} are close on $\overline{\Omega} \setminus Q_i^\xi$, and the intrinsic metric of Y_i increase on Q_i^ξ with respect to Y_{i-1} . From this, we obtain that the norm of Y_i is greater than $r + s$ around Q_i . To do this, we use the López-Ros transformation in the set of Cartesian coordinates T_i and Runge's theorem. So, note that the third coordinates of $X_n(a_i)$ and $X_n(a_{i+1})$, in the frame T_i , are close and greater than $r + s$. This fact will be decisive in Section 4.3.

Previously, we define a collection of neighborhoods of the arcs C_i and a collection of neighborhoods of the curves Q_i :

For all $i = 1 \dots, n$, let $\widehat{C}_i$ be an open neighborhood of C_i , small enough to satisfy:

$$\|X_n(z) - X_n(a_i)\| < 3\epsilon_0, \quad \forall z \in \widehat{C}_i \cap \overline{\Omega} \quad (23)$$

We can take $\widehat{C}_i$ verifying the above because, using $(A4)_i$, for all $z \in C_i$, $\|X_n(z) - X_n(a_i)\| < \|X_i(z) - X_i(a_i)\| + 2\epsilon_0 < 3\epsilon_0$.

We define $0 < \epsilon_1$ such that

$$2\epsilon_1 < \frac{1}{2} \min_{Q_i} \{|f_{(X_n, T_i)}|\}, \quad \forall i = 1, \dots, n. \quad (24)$$

Observe that from 6.f), $\min_{Q_i} \{|f_{(X_n, T_i)}|\} \neq 0$.

For all $i = 1, \dots, n$, we define $Q_i^\xi = \{z \in \mathbb{C} : \operatorname{dist}(z, Q_i) \leq \xi\}$, where $\operatorname{dist}(z, A)$ means the Euclidean distance between point z and set A . We assume that $0 < \xi$ is small enough to satisfy:

7.a) $Q_i^\xi \subset \Omega'$;

7.b) $Q_i^\xi \cap Q_j^\xi = \emptyset$, $i \neq j$;




Figure 4: Properties of X_n .

7.c) $Q_i^\xi \cap \overline{D(p_k, \delta)} = \emptyset$, for $k \notin \{i, i+1\}$;

7.d) $Q_i^\xi \subset B^i$;

7.e) $Q_i^{\xi/2}$ and $\overline{\Omega \setminus Q_i^\xi}$ are simply connected sets;

7.f) For all $z \in Q_i$ we have $|f_{(X_n, T_i)}(z) - f_{(X_n, T_i)}(x)| < \epsilon_1$, $\forall x \in B(z, \xi/2)$;

7.g) $\sup_{z \in \Omega \setminus Q_i^\xi} \inf \{\text{Euclidean length of } \alpha : \alpha \text{ is a curve in } \overline{\Omega \setminus Q_i^\xi} \text{ with origin 0 and ending at } z\} < l$.

Label $Y_0 \stackrel{\text{def}}{=} X_n$. We are going to construct in a recursive way $Y_1, \dots, Y_n$, where $Y_i : \Omega' \rightarrow \mathbb{R}^3$ is a conformal minimal immersion with $Y_i(0) = 0$ verifying the following properties:

(B1)_i $(Y_i)_{3, T_i} = (Y_{i-1})_{3, T_i}$;

(B2)_i $\|Y_i(z) - Y_{i-1}(z)\| < \frac{\epsilon_0}{n}$, $\forall z \in \overline{\Omega \setminus Q_i^\xi}$;

(B3)_i $|f_{(Y_i, T_k)}(z) - f_{(Y_{i-1}, T_k)}(z)| < \frac{\epsilon_1}{n}$, $\forall z \in \overline{\Omega \setminus Q_i^\xi}$, for $k = i+1, \dots, n$;

(B4)_i $\frac{\xi}{2} \left(\left(\frac{1}{\tau_i} + \frac{\nu_i}{\tau_i(\tau_i - \nu_i)} \right) \max_{Q_i^\xi} \{|f_{(Y_{i-1}, T_i)} g_{(Y_{i-1}, T_i)}^2|\} + \nu_i \max_{Q_i^\xi} \{|f_{(Y_{i-1}, T_i)}|\} \right) < 1$;

(B5)_i $\frac{1}{2} \left(\frac{\tau_i \xi}{4} \min_{Q_i} \{|f_{(Y_0, T_i)}|\} - 1 \right) > 2(r+s) + 1$;

where τ_i and ν_i are certain positive constants.

Assume we have $Y_0, Y_1, \dots, Y_{i-1}$. We define Y_i as:

$$f_{(Y_i, T_i)} = f_{(Y_{i-1}, T_i)} l_i, \quad g_{(Y_i, T_i)} = g_{(Y_{i-1}, T_i)} / l_i,$$

where $l_i : \mathbb{C} \rightarrow \mathbb{C}$ is a holomorphic non zero function verifying:

- $|l_i(z) - \tau_i| < \nu_i$, $\forall z \in Q_i^{\xi/2}$;
- $|l_i(z) - 1| < \nu_i$, $\forall z \in \overline{\Omega \setminus Q_i^\xi}$.

The existence of this function [4] is a consequence of Runge's theorem. We define Y_i as the minimal immersion that becomes from the above meromorphic data.

We have assumed that $\tau_i > 0$ is large enough and $\nu_i > 0$ is small enough to satisfy (B2)_i, (B3)_i, (B4)_i and (B5)_i. It is possible because the Weierstrass representation of Y_{i-1} uniformly converges to the Weierstrass representation of Y_i , as $\nu_i \rightarrow 0$ and $\tau_i \rightarrow \infty$ on $\overline{\Omega \setminus Q_i^\xi}$. We take into account Property 7.g), to obtain (B2)_i.

Observe that (B1)_i trivially holds.

4.3 Definition of Y

Eventually, we define the open set U as $U = \Omega$ and $Y : U \rightarrow \mathbb{R}^3$ as $Y = Y_n$. In this section, we are going to check that the immersion Y verifies all the claims of the lemma.

Bearing in mind $\overline{\text{Int } P} \subset \Omega \setminus ((\cup_{k=1}^n D(p_k, \delta)) \cup (\cup_{k=1}^n Q_k^\xi))$, Property (b) is a easy consequence of (P1) and (B2).

The proof of the rest of claims of Lemma are nontrivial:

Properties (a) and (c): To prove the existence of a polygon Q verifying (a) and (c), we only need to get the following:

$$\begin{aligned} & \text{for all curve } \beta \text{ with origin } 0 \text{ and ending at a} \\ & \text{point of } \partial\Omega, \text{ there exists } z' \in \beta \text{ such that } \|Y(z')\| > r + s. \end{aligned} \quad (25)$$

We are going to see the above property. Let $\beta \subset \overline{\Omega}$ be a curve with $\beta(0) = 0$ and $\beta(1) = z_0 \in \partial\Omega$.

First, we want to show that $\|Y_n(z)\| > r + s$, $\forall z \in \widehat{C}_i$, for $i = 1, \dots, n$. Let $z \in \widehat{C}_i$, we study three possibilities:

- Suppose that $z \in \widehat{C}_i \cap Q_i^\xi$, $i \in \{1, \dots, n\}$. Then, considering T_i , and using (B2), (B1)_i, (23), (22) and (P4)_i, we have

$$\begin{aligned} \|Y_n(z)\| &\geq \|Y_i(z)\| - \epsilon_0 \geq |(Y_i(z))_3| - \epsilon_0 = |(Y_{i-1}(z))_3| - \epsilon_0 \geq |(Y_0(z))_3| - 2\epsilon_0 > \\ &> |(Y_0(a_i))_3| - 5\epsilon_0 = \|Y_0(a_i)\| - 5\epsilon_0 > \lambda(r + s) - 5\epsilon_0 > r + s. \end{aligned}$$

- Suppose that $z \in \widehat{C}_i \cap Q_{i-1}^\xi$. Now, considering T_{i-1} , one has

$$\|Y_n(z)\| \geq |(Y_0(z))_{3, T_{i-1}}| - 2\epsilon_0 \geq$$

and from (23), (P3)_{i-1} and (P4)_{i-1}, we deduce

$$\begin{aligned} &\geq |(Y_0(a_i))_{3, T_{i-1}}| - 5\epsilon_0 \geq |(Y_0(a_{i-1}))_{3, T_{i-1}}| - \epsilon_0(4\lambda^2 s' + 17) - 5\epsilon_0 \geq \\ &\geq \lambda(r + s) - \epsilon_0(4\lambda^2 s' + 17) - 5\epsilon_0 \geq r + s. \end{aligned}$$

Note that case $\widehat{C}_1 \cap Q_n^\xi$ can be proved in the same way.

- Suppose that $z \in \widehat{C}_i \setminus \cup_{k=1}^n Q_k^\xi$. Therefore,

$$\|Y_n(z)\| \geq \|Y_0(z)\| - \epsilon_0 > \|Y_0(a_i)\| - 4\epsilon_0 > \lambda(r + s) - 4\epsilon_0 > r + s.$$

This proves that $\|Y_n(z)\| > r+s, \forall z \in \widehat{C}_i$. Hence, we can suppose that $\beta \cap \widehat{C}_i = \emptyset$, $\forall i = 1, \dots, n$.

From 6.c), $z_0 \in Q_i$, $i = 1, \dots, n$. Let $z_1 \in \beta \cap \partial B(z_0, \xi/2)$. Observe that $\overline{z_1 z_0} \subset Q_i^{\xi/2}$. Next, we give a lower bound for $\|Y_n(z_0) - Y_n(z_1)\|$. We consider T_i , and write $f^{i-1} = f_{(Y_{i-1}, T_i)}$, $g^{i-1} = g_{(Y_{i-1}, T_i)}$. Then,

$$\begin{aligned}
\|Y_n(z_0) - Y_n(z_1)\| &\geq \|Y_i(z_0) - Y_i(z_1)\| - 2\epsilon_0 \geq \\
&\geq \left| \left(\operatorname{Re} \frac{1}{2} \int_{\overline{z_1 z_0}} f^{i-1} l_i - \frac{f^{i-1}(g^{i-1})^2}{l_i} \right) + \right. \\
&\quad \left. + i \left(\operatorname{Re} \frac{i}{2} \int_{\overline{z_1 z_0}} f^{i-1} l_i + \frac{f^{i-1}(g^{i-1})^2}{l_i} \right) \right| - 2\epsilon_0 = \\
&= \frac{1}{2} \left| \int_{\overline{z_1 z_0}} \overline{f^{i-1} l_i} - \frac{f^{i-1}(g^{i-1})^2}{l_i} \right| - 2\epsilon_0 \geq \\
&\geq \frac{\tau_i}{2} \left| \int_{\overline{z_1 z_0}} \overline{f^{i-1}} \right| - \frac{1}{2\tau_i} \left| \int_{\overline{z_1 z_0}} f^{i-1}(g^{i-1})^2 \right| - \\
&\quad - \frac{1}{2} \left| \int_{\overline{z_1 z_0}} \overline{f^{i-1}(l_i - \tau_i)} \right| - \frac{1}{2} \left| \int_{\overline{z_1 z_0}} f^{i-1}(g^{i-1})^2 \left(\frac{1}{l_i} - \frac{1}{\tau_i} \right) \right| - 2\epsilon_0 \geq \\
&\geq \frac{\tau_i}{2} \left| \int_{\overline{z_1 z_0}} \overline{f^{i-1}} \right| - \frac{\xi}{4} \left(\frac{1}{\tau_i} \max_{Q_i^\xi} \{|f^{i-1}(g^{i-1})^2|\} + \right. \\
&\quad \left. + \nu_i \max_{Q_i^\xi} \{|f^{i-1}|\} + \frac{\nu_i}{\tau_i(\tau_i - \nu_i)} \max_{Q_i^\xi} \{|f^{i-1}(g^{i-1})^2|\} \right) - 2\epsilon_0 \geq
\end{aligned}$$

using (B4)_i, we continue

$$\geq \frac{1}{2} \left(\tau_i \left| \int_{\overline{z_1 z_0}} \overline{f^{i-1}} \right| - 1 \right) - 2\epsilon_0.$$

On the other hand, from (B3)₁, ..., (B3)_{i-1}, 7.f), and (24), we deduce

$$\begin{aligned}
\left| \int_{\overline{z_1 z_0}} \overline{f^{i-1}(w)} dw \right| &\geq \\
&\geq \left| f_{(Y_0, T_i)}(z_0) \int_{\overline{z_1 z_0}} dw \right| - \left| \int_{\overline{z_1 z_0}} (f_{(Y_0, T_i)}(z_0) - f_{(Y_0, T_i)}(w)) dw \right| - \\
&\quad - \left| \int_{\overline{z_1 z_0}} (f_{(Y_0, T_i)}(w) - f^{i-1}(w)) dw \right| \geq \\
&\geq \frac{\xi}{2} (|f_{(Y_0, T_i)}(z_0)| - \epsilon_1 - \epsilon_1) \geq \frac{\xi}{2} (\min_{Q_i} \{|f_{(Y_0, T_i)}|\} - 2\epsilon_1) \geq \\
&\geq \frac{\xi}{4} \min_{Q_i} \{|f_{(Y_0, T_i)}|\}.
\end{aligned}$$

Therefore,

$$\|Y_n(z_0) - Y_n(z_1)\| \geq \frac{1}{2} \left(\tau_i \frac{\xi}{4} \min_{Q_i} \{|f_{(Y_0, T_i)}|\} - 1 \right) - 2\epsilon_0 \geq$$

and using (B5)_i, we obtain

$$\|Y_n(z_0) - Y_n(z_1)\| \geq 2(r + s) + 1 - 2\epsilon_0 > 2(r + s). \quad (26)$$

Indeed, one has $\|Y_n(z_0)\| > r + s$ or $\|Y_n(z_1)\| > r + s$.

This conclude the proof of (25). Thus, there is a polygon Q verify (a) and (c).

Property (d): Let $z \in \text{Int } Q \setminus \text{Int } P$. We distinguish the five possible cases to prove this property:

- Suppose that $z \notin (\cup_{k=1}^n D(p_i, \delta)) \cup (\cup_{k=1}^n Q_k^\xi)$. (P1), (B2) and (1) imply $\|Y_n(z)\| \geq \|X(z)\| - 2\epsilon_0 > r - 2\epsilon_0 > r - 3\sqrt{sr}$.
- Suppose that $z \in D(p_i, \delta) \setminus \cup_{k=1}^n Q_k^\xi$. We consider $S(p_i)$ and use (P1), (B2), (P2)_i, then:

$$\begin{aligned} \|Y_n(z)\| &\geq \|X_i(z)\| - 2\epsilon_0 \geq |(X_i(z))_3| - 2\epsilon_0 = \\ &= |(X_{i-1}(z))_3| - 2\epsilon_0 \geq |(X(z))_3| - 3\epsilon_0 \geq \end{aligned}$$

and, from (4) and the definition of $S(p_i)$, one has

$$\geq |(X(p_i))_3| - 4\epsilon_0 = \|X(p_i)\| - 4\epsilon_0 > r - 4\epsilon_0 > r - 3\sqrt{sr}.$$

- Suppose that $z \in D(p_i, \delta) \cap Q_i^\xi$. Now, we consider the orthogonal frame T_i , and obtain

$$\begin{aligned} \|Y_n(z)\| &\geq \|Y_i(z)\| - \epsilon_0 \geq |(Y_i(z))_3| - \epsilon_0 = |(Y_{i-1}(z))_3| - \epsilon_0 \geq |(Y_0(z))_3| - 2\epsilon_0 \geq \\ &\geq |(Y_0(a_i))_3| - |(Y_0(a_i))_3 - (Y_0(q_i))_3| - |(Y_0(q_i))_3 - (Y_0(z))_3| - 2\epsilon_0 > \end{aligned}$$

Note that $|(Y_0(a_i))_{3, T_i}| = \|Y_0(a_i)\|$. Since $z \in G_i$, we can use (P4)_i and (P5)_i to get the following

$$> \lambda(r + s) - 2\lambda^3 s' - 12\epsilon_0 = \lambda(r + s) - 2\lambda^3 \sqrt{\lambda^4(r + s)^2 - r^2} - 12\epsilon_0 >$$

next we recall the definition of λ on 1.b), and $s < r/100$, then

$$> r - 2\sqrt{(r + s)^2 - r^2} - 12\epsilon_0 > r - 3\sqrt{sr}.$$

- Suppose that $z \in D(p_{i+1}, \delta) \cap Q_i^\xi$. We consider T_i . This case is similar to the former case:

$$\begin{aligned} \|Y_n(z)\| &\geq |(Y_0(a_i))_{3,T_i}| - |(Y_0(a_i))_{3,T_i} - (Y_0(a_{i+1}))_{3,T_i}| - \\ &\quad - |(Y_0(a_{i+1}))_{3,T_i} - (Y_0(q_{i+1}))_{3,T_i}| - |(Y_0(q_{i+1}))_{3,T_i} - (Y_0(z))_{3,T_i}| - 2\epsilon_0 \geq \end{aligned}$$

Here, $(P3)_i$ gives

$$> \lambda(r + s) - \epsilon_0(4\lambda^2 + 17) - 2\lambda^3 s' - 12\epsilon_0 \geq r - 3\sqrt{sr}.$$

Note that case $z \in D(p_1, \delta) \cap Q_n^\xi$ can be proved in the same way, considering T_n .

- Suppose that $z \in Q_i^\xi \setminus \cup_{k=1}^n D(p_k, \delta)$. We consider T_i . A similar computation gives

$$\begin{aligned} \|Y_n(z)\| &\geq |(Y_0(a_i))_3| - |(Y_0(a_i))_3 - (Y_0(q_i))_3| - |(Y_0(q_i))_3 - (Y_0(z))_3| - 2\epsilon_0 \geq \\ &\geq \lambda(r + s) - \lambda^3 s' - 7\epsilon_0 - |(Y_0(q_i))_3 - (Y_0(z))_3| \end{aligned}$$

and using properties (P1) and (4), we obtain

$$\geq \lambda(r + s) - \lambda^3 s' - 10\epsilon_0 \geq r - 3\sqrt{sr}.$$

This completes the proof of the lemma.

References

- [1] P. Collin, R. Kusner, W. H. Meeks III, and H. Rosenberg *The topology, geometry and conformal structures of properly embedded minimal surfaces*. In preparation.
- [2] F. Martín and S. Morales, *A note about the asymptotic behavior of a complete bounded minimal surface in $\mathbb{R}^3$* . Preprint.
- [3] W. H. Meeks III, *Global problems in Classical minimal surfaces theory*. In preparation.
- [4] N. Nadirashvili, *Hadamard's and Calabi-Yau's conjectures on negatively curved and minimal surfaces*. Invent. math., 126 (1996), 457-465.

Santiago Morales

Departamento de Geometría y Topología

Universidad de Granada

18071 Granada, Spain

E-mail: santimo@ugr.es

URL: <http://www.ugr.es/~santimo>